

LOW-DIMENSIONAL HOMOLOGY OF $SL_2(k[t, t^{-1}])$

KEVIN HUTCHINSON

ABSTRACT. We prove analogues of the fundamental theorem of K -theory for the second and third homology of SL_2 over an infinite field. The statements of the theorems involve Milnor-Witt K -theory and refined scissors congruence groups. We use these results to calculate the low-dimensional homology of SL_2 of Laurent polynomials over certain fields.

1. INTRODUCTION

Our goal in this paper is to prove unstable analogues of the fundamental theorem of K -theory for the second and third homology of SL_2 over an infinite field k , and to use these results to calculate the low-dimensional homology of SL_2 of Laurent polynomials over certain fields.

We follow the approach of K. Knudson ([7]) who studied the homology of $SL_2(k[t, t^{-1}])$ via the Mayer-Vietoris sequence associated its natural decomposition as an amalgamated product.

Let k be an infinite field. When $n \geq 3$ then $H_2(SL_n(k), \mathbb{Z}) \cong K_2(k)$. Furthermore, by [8, Corollary 6.11]

$$H_2(SL_n(k[t, t^{-1}]), \mathbb{Z}) \cong K_2(k[t, t^{-1}]) \text{ for } n \geq 3.$$

The fundamental theorem of K -theory, combined with the fact that $K_n(A) \cong K_n(A[t])$ for regular rings A ([16]), tells us that

$$K_2(k[t, t^{-1}]) \cong K_2(k) \oplus K_1(k).$$

Here the homomorphism $K_2(k[t, t^{-1}]) \rightarrow K_1(k)$ is the composite

$$K_2(k[t, t^{-1}]) \longrightarrow K_2(k(t)) \xrightarrow{\delta_t} K_1(k)$$

where δ_t is the tame symbol associated to the t -adic valuation on $k(t)$.

It follows that

$$H_2(SL_n(k[t, t^{-1}]), \mathbb{Z}) \cong H_2(SL_n(k), \mathbb{Z}) \oplus K_1(k)$$

for all $n \geq 3$.

In section 4 (Theorem 4.1) below we prove an analogue of this for the case $n = 2$: Let k be an infinite field of characteristic not equal to 2. Then

$$H_2(SL_2(k[t, t^{-1}]), \mathbb{Z}) \cong H_2(SL_2(k), \mathbb{Z}) \oplus K_1^{\text{MW}}(k).$$

Here $K_1^{\text{MW}}(k)$ is the first Milnor-Witt K -theory group of the field k (see section 3). There is a natural surjective map $K_1^{\text{MW}}(k) \rightarrow K_1(k) = k^\times$ whose kernel is the second power of the fundamental ideal in the Witt ring of k . Furthermore, for any infinite field K there is a natural isomorphism $H_2(SL_2(k), \mathbb{Z}) \cong K_2^{\text{MW}}(K)$ and the homomorphism $H_2(SL_2(k[t, t^{-1}]), \mathbb{Z}) \rightarrow K_1^{\text{MW}}(k)$ is the composite

$$H_2(SL_2(k[t, t^{-1}]), \mathbb{Z}) \longrightarrow H_2(SL_2(k(t)), \mathbb{Z}) \cong K_2^{\text{MW}}(k(t)) \xrightarrow{\delta_t} K_1^{\text{MW}}(k)$$

Date: April 24, 2014.

1991 Mathematics Subject Classification. 19G99, 20G10.

Key words and phrases. K -theory, Group Homology .

where δ_t is a connecting homomorphism, analogous to the tame symbol, arising in Milnor-Witt K -theory.

The third homology of the special linear groups of fields is also closely related to K -theory. Combining [19, Corollary 5.2] and [6, Theorem 4.7] it follows that

$$H_3(\mathrm{SL}_n(k), \mathbb{Z}) \cong K_3(k)/(\{-1\} \cdot K_2(k)) \text{ for all } n \geq 3.$$

In particular,

$$H_3(\mathrm{SL}_n(k), \mathbb{Z}[\frac{1}{2}]) \cong K_3(k) \otimes \mathbb{Z}[\frac{1}{2}]$$

for all $n \geq 3$.

By contrast, when $n = 2$, the relation between $H_3(\mathrm{SL}_2(k), \mathbb{Z})$ and $K_3(k)$ is much more remote. There is a natural surjective map from $H_3(\mathrm{SL}_2(k), \mathbb{Z})$ to $K_3^{\mathrm{ind}}(k)$ but for general infinite fields the kernel may be very large (see [4, Theorem 5.1]).

However, recent computations by the author suggest a natural candidate for the functor

$$\frac{H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z})}{H_3(\mathrm{SL}_2(k), \mathbb{Z})},$$

at least when replace $\mathbb{Z}$ by $\mathbb{Z}[\frac{1}{2}]$: Namely, given a field K with a discrete valuation v and residue field k there is a natural homomorphism $\delta_\pi : H_2(\mathrm{SL}_2(K), \mathbb{Z}[\frac{1}{2}]) \rightarrow \mathcal{RP}_1(k) \otimes \mathbb{Z}[\frac{1}{2}]$ associated to a uniformizer π . Here $\mathcal{RP}_1(k)$ is a *refined scissors congruence group* (see section 6 below). These groups arise naturally in the calculation of the third homology of SL_2 of fields and local rings.

In [3], the author has shown that for certain families of discretely valued fields K with residue field k and valuation ring $\mathcal{O}$ there is a natural exact localization sequence

$$0 \rightarrow H_3(\mathrm{SL}_2(\mathcal{O}), \mathbb{Z}[\frac{1}{2}]) \rightarrow H_3(\mathrm{SL}_2(K), \mathbb{Z}[\frac{1}{2}]) \rightarrow \mathcal{RP}_1(k) \otimes \mathbb{Z}[\frac{1}{2}] \rightarrow 0.$$

The main result of section 7 below is that for any infinite field k there is a natural isomorphism

$$H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \cong H_3(\mathrm{SL}_2(k), \mathbb{Z}[\frac{1}{2}]) \oplus (\mathcal{RP}_1(k) \otimes \mathbb{Z}[\frac{1}{2}]).$$

Here the map $H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \rightarrow \mathcal{RP}_1(k) \otimes \mathbb{Z}[\frac{1}{2}]$ is the composite

$$H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \longrightarrow H_3(\mathrm{SL}_2(k(t)), \mathbb{Z}[\frac{1}{2}]) \xrightarrow{\delta_t} \mathcal{RP}_1(k) \otimes \mathbb{Z}[\frac{1}{2}].$$

A key ingredient in our calculations is Knudson's homotopy invariance theorem ([8, Theorem 3.4]): If k is an infinite field, then the inclusion $\mathrm{SL}_2(k) \rightarrow \mathrm{SL}_2(k[t])$ induces an isomorphism on integral homology

$$H_\bullet(\mathrm{SL}_2(k), \mathbb{Z}) \cong H_\bullet(\mathrm{SL}_2(k[t]), \mathbb{Z}).$$

However, the example of Krstic and McCool ([10]) shows that even for H_1 this result does not extend to more general ground rings k . In particular, the corresponding statement for the ring of polynomials in two or more variables is not true.

1.1. Preliminaries and notation. For a commutative ring A the short exact sequence

$$1 \rightarrow \mathrm{SL}_2(A) \rightarrow \mathrm{GL}_2(A) \rightarrow A^\times \rightarrow 1$$

gives rise to a natural action of $A^\times$ on $H_\bullet(\mathrm{SL}_2(A), \mathbb{Z})$: given $a \in A^\times$, choose $M = M(a) \in \mathrm{GL}_2(A)$ with determinant a . Conjugation by M is an automorphism of $\mathrm{SL}_2(A)$ whose induced action on $H_\bullet(\mathrm{SL}_2(A), \mathbb{Z})$ depends only on a and not on the choice of M . Furthermore, since a^2 is the determinant of a scalar matrix, the action of a^2 on $H_\bullet(\mathrm{SL}_2(A), \mathbb{Z})$ is trivial for any

$a \in A^\times$. It follows that each of the groups $H_n(\mathrm{SL}_2(A), \mathbb{Z})$ is naturally a module over the group ring $\mathbb{Z}[A^\times/(A^\times)^2]$ of square classes of units of A .

For $a \in A^\times$, the square class of a will be denoted $\langle a \rangle \in \mathbb{Z}[A^\times/(A^\times)^2]$. Furthermore, the element $\langle a \rangle - 1$ in the augmentation ideal, $\mathcal{I}_A$, will be denoted $\langle\langle a \rangle\rangle$.

For an abelian group M we will let $M\left[\frac{1}{2}\right]$ denote the $\mathbb{Z}\left[\frac{1}{2}\right]$ -module $M \otimes \mathbb{Z}\left[\frac{1}{2}\right]$.

2. THE MAYER-VIETORIS SEQUENCE

Let k be an infinite field and let $G = \mathrm{SL}_2(k[t, t^{-1}])$. We denote the matrix

$$\begin{bmatrix} t & 0 \\ 0 & 1 \end{bmatrix} \in \mathrm{GL}_2(k[t, t^{-1}]) \subset \mathrm{GL}_2(k(t))$$

by $A(t)$. $A(t)$ normalizes $\mathrm{SL}_2(k(t))$ and $\mathrm{SL}_2(k[t, t^{-1}])$. We denote by C_t the conjugation map $g \mapsto A(t)^{-1}gA(t) = g^{A(t)}$ on either of these subgroups. Observe, in particular, that the action of $A(t)$ by conjugation on G induces on $H_\bullet(G, \mathbb{Z})$ multiplication by $\langle t \rangle \in \mathbb{Z}[k[t, t^{-1}]^\times/(k[t, t^{-1}]^\times)^2]$.

Let G_1 denote the subgroup $\mathrm{SL}_2(k[t])$ of G and let

$$G_2 := C_t(G_1) = \left\{ \begin{bmatrix} a & t^{-1}b \\ tc & d \end{bmatrix} \in G : a, b, c, d \in k[t], ad - bc = 1 \right\}.$$

Let

$$\Gamma = G_1 \cap G_2 = \left\{ \begin{bmatrix} a & b \\ tc & d \end{bmatrix} \in G : a, b, c, d \in k[t], ad - tbc = 1 \right\}.$$

K. Knudson ([7]) observes that since $k[t, t^{-1}]$ is a dense subring of the complete discretely valued field $k((t))$, Serre's theory of trees (see [17, Chapter II, Theorem 3]) allows us to deduce:

Theorem 2.1. *$G = \mathrm{SL}_2(k[t, t^{-1}])$ is the sum of the subgroups G_1 and G_2 amalgamated along their common intersection Γ :*

$$G = \mathrm{SL}_2(k[t, t^{-1}]) = \mathrm{SL}_2(k[t]) \star_\Gamma \mathrm{SL}_2(k[t])^{A(t)} = G_1 \star_\Gamma G_2.$$

This amalgamated product decomposition gives us immediately a short exact sequence of $\mathbb{Z}[G]$ -modules

$$0 \longrightarrow \mathbb{Z}[G/\Gamma] \xrightarrow{\alpha} \mathbb{Z}[G/G_1] \oplus \mathbb{Z}[G/G_2] \xrightarrow{\beta} \mathbb{Z} \longrightarrow 0.$$

where α is the map $g\Gamma \mapsto (gG_1, gG_2)$ and β is the unique $\mathbb{Z}[G]$ -homomorphism sending $(G_1, 0)$ to -1 and $(0, G_2)$ to 1 .

The associated long exact sequence in homology – the Mayer-Vietoris sequence of the amalgamated product – takes the form

$$\cdots \xrightarrow{\delta} H_i(\Gamma, \mathbb{Z}) \xrightarrow{\alpha} H_i(G_1, \mathbb{Z}) \oplus H_i(G_2, \mathbb{Z}) \xrightarrow{\beta} H_i(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \xrightarrow{\delta} \cdots$$

Implicit here are the isomorphisms of Shapiro's Lemma: If H is a subgroup of K , then the inclusion $\mathbb{Z} \rightarrow \mathbb{Z}[G/H]$ of $\mathbb{Z}[H]$ -modules induces an isomorphism $H_\bullet(H, \mathbb{Z}) \cong H_\bullet(G, \mathbb{Z}[G/H])$.

Known results allow us to simplify further some terms in this sequence. To begin with we have the following homotopy-invariance property of the homology of $\mathrm{SL}_2(k)$ (see [9, Theorem 4.3.1]):

Theorem 2.2. *Let k be an infinite field. Then the inclusion $k \rightarrow k[t]$ induces an isomorphism on integral homology*

$$H_\bullet(\mathrm{SL}_2(k), \mathbb{Z}) \cong H_\bullet(\mathrm{SL}_2(k[t]), \mathbb{Z}).$$

We let $\iota : k^\times \rightarrow \mathrm{SL}_2(k)$ denote the embedding

$$a \mapsto \begin{bmatrix} a & 0 \\ 0 & a^{-1} \end{bmatrix}.$$

The image of ι lies in the subgroup

$$\Gamma \cap \mathrm{SL}_2(k) = B(k) := \left\{ \begin{bmatrix} a & b \\ 0 & a^{-1} \end{bmatrix} : a \in k^\times, b \in k \right\}.$$

Furthermore, we have ([9, Theorem 4.4.1])

Theorem 2.3. *When the field k is infinite, ι induces isomorphisms on integral homology*

$$H_\bullet(k^\times, \mathbb{Z}) \cong H_\bullet(B, \mathbb{Z}) \cong H_\bullet(\Gamma, \mathbb{Z}).$$

With all of these identifications, together with the fact that C_t induces an isomorphism from G_1 to G_2 , the Mayer-Vietoris sequence takes the form

$$\cdots \xrightarrow{\delta} H_i(k^\times, \mathbb{Z}) \xrightarrow{\alpha} H_i(\mathrm{SL}_2(k), \mathbb{Z}) \oplus H_i(\mathrm{SL}_2(k), \mathbb{Z}) \xrightarrow{\beta} H_i(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \xrightarrow{\delta} \cdots$$

where $\alpha(z) = (\iota(z), \iota(z))$ and $\beta(z_1, z_2) = \langle t \rangle j(z_2) - j(z_1)$.

As Knudson points out, the existence of this exact sequence immediately implies that

$$H_1(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) = 0$$

for any infinite field k . (This fact is originally due to P. Cohn, [1].)

3. MILNOR-WITT K -THEORY AND H_2 OF SL_2

In this section all fields are of characteristic different from 2.

The theorem of Matusmoto and Moore gives an explicit presentation of $H_2(\mathrm{SL}_2(k), \mathbb{Z})$ for an infinite field k . In [18], A. Suslin showed that this implies a natural isomorphism

$$(1) \quad H_2(\mathrm{SL}_2(k), \mathbb{Z}) \cong I^2(k) \times_{I^2(k)/I^3(k)} K_2^M(k)$$

where $I^n(k)$ denotes the n -th power of the fundamental ideal of the Witt ring $W(k)$ of the field k .

We can now recognize the group on the right-hand side of this isomorphism as the second *Milnor-Witt K -group* of the field k . The Milnor-Witt K -theory of a field is the graded ring $K_\bullet^{\mathrm{MW}}(F)$ with the following presentation (due to F. Morel and M. Hopkins, see [12]):

Generators: $[a]$, $a \in F^\times$, in dimension 1 and a further generator η in dimension -1 .

Relations:

- (a) $[ab] = [a] + [b] + \eta \cdot [a] \cdot [b]$ for all $a, b \in F^\times$
- (b) $[a] \cdot [1 - a] = 0$ for all $a \in F^\times \setminus \{1\}$
- (c) $\eta \cdot [a] = [a] \cdot \eta$ for all $a \in F^\times$
- (d) $\eta \cdot h = 0$, where $h = \eta \cdot [-1] + 2 \in K_0^{\mathrm{MW}}(F)$.

Clearly there is a unique surjective homomorphism of graded rings $K_\bullet^{\mathrm{MW}}(F) \rightarrow K_\bullet^M(F)$ sending $[a]$ to $\{a\}$ and inducing an isomorphism

$$\frac{K_\bullet^{\mathrm{MW}}(F)}{\langle \eta \rangle} \cong K_\bullet^M(F)$$

(where $K_n^M(F) := 0$ for $n < 0$).

Furthermore, there is a natural surjective homomorphism of graded rings $K_{\bullet}^{\text{MW}}(F) \rightarrow I^{\bullet}(F)$ sending $[a]$ to $\langle\langle a \rangle\rangle$ and η to η . Morel shows that there is an induced isomorphism of graded rings

$$\frac{K_{\bullet}^{\text{MW}}(F)}{\langle h \rangle} \cong I^{\bullet}(F)$$

(where $I^n(F) := W(F)$ for $n < 0$).

The main structure theorem on Milnor-Witt K -theory is the following theorem of Morel:

Theorem 3.1 (Morel, [13]). *For a field F of characteristic not 2, the commutative square of graded rings*

$$\begin{array}{ccc} K_{\bullet}^{\text{MW}}(F) & \longrightarrow & K_{\bullet}^{\text{M}}(F) \\ \downarrow & & \downarrow \\ I^{\bullet}(F) & \longrightarrow & I^{\bullet}(F)/I^{\bullet+1}(F) \end{array}$$

is cartesian.

Thus for each $n \in \mathbb{Z}$ we have an isomorphism

$$(2) \quad K_n^{\text{MW}}(F) \cong K_n^{\text{M}}(F) \times_{I^n(F)/I^{n+1}(F)} I^n(F).$$

It follows that for all n there is a natural short exact sequence

$$(3) \quad 0 \rightarrow I^{n+1}(F) \rightarrow K_n^{\text{MW}}(F) \rightarrow K_n^{\text{M}}(F) \rightarrow 0$$

where the inclusion $I^{n+1}(F) \rightarrow K_n^{\text{MW}}(F)$ is given by

$$\langle\langle a_1, \dots, a_{n+1} \rangle\rangle \mapsto \eta[a_1] \cdots [a_{n+1}].$$

Similarly, for $n \geq 0$, there is a short exact sequence

$$0 \rightarrow 2K_n^{\text{M}}(F) \rightarrow K_n^{\text{MW}}(F) \rightarrow I^n(F) \rightarrow 0$$

where the inclusion $2K_n^{\text{M}}(F) \rightarrow K_n^{\text{MW}}(F)$ is given (for $n \geq 1$) by

$$2\{a_1, \dots, a_n\} \mapsto h[a_1] \cdots [a_n].$$

For $n \geq 2$ we have

$$h[a_1][a_2] \cdots [a_n] = ([a_1][a_2] - [a_2][a_1])[a_3] \cdots [a_n] = [a_1^2][a_2] \cdots [a_n].$$

In particular, by (1) and (2), there is a natural isomorphism

$$H_2(\text{SL}_2(k), \mathbb{Z}) \cong K_2^{\text{MW}}(k).$$

Indeed, the resulting diagram

$$\begin{array}{ccc} H_2(\text{SL}_2(k), \mathbb{Z}) & \xrightarrow{\cong} & K_2^{\text{MW}}(k) \\ \downarrow & & \downarrow \\ H_2(\text{SL}_3(k), \mathbb{Z}) & \xrightarrow{\cong} & K_2^{\text{M}}(k) = K_2(k) \end{array}$$

commutes.

4. THE SECOND HOMOLOGY OF $\mathrm{SL}_2(k[t, t^{-1}])$

In this section k is an infinite field of characteristic not equal to 2.

Our goal is to prove:

Theorem 4.1. *Let k be an infinite field of characteristic not equal to 2. Then*

$$H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \cong H_2(\mathrm{SL}_2(k), \mathbb{Z}) \oplus K_1^{\mathrm{MW}}(k).$$

The Mayer-Vietoris sequence gives us an exact sequence

$$H_2(k^\times, \mathbb{Z}) \longrightarrow H_2(\mathrm{SL}_2(k), \mathbb{Z}) \oplus H_2(\mathrm{SL}_2(k), \mathbb{Z}) \longrightarrow H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \xrightarrow{\delta} H_1(k^\times, \mathbb{Z}) \longrightarrow 0.$$

However, since the homomorphism $j : H_i(\mathrm{SL}_2(k), \mathbb{Z}) \rightarrow H_i(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z})$ is a split injection, we obtain an induced exact sequence

$$H_2(k^\times, \mathbb{Z}) \xrightarrow{\iota} H_2(\mathrm{SL}_2(k), \mathbb{Z}) \xrightarrow{\bar{\beta}} \frac{H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z})}{H_2(\mathrm{SL}_2(k), \mathbb{Z})} \xrightarrow{\delta} H_1(k^\times, \mathbb{Z}) \longrightarrow 0$$

where $\bar{\beta}(z) = \langle t \rangle j(z)$.

Recall that there is a natural isomorphism $K_2^{\mathrm{MW}}(k) \cong H_2(\mathrm{SL}_2(k), \mathbb{Z})$. Given $a, b \in k^\times$, we will let $[a, b]$ denote the element of $H_2(\mathrm{SL}_2(k), \mathbb{Z})$ corresponding, under this isomorphism, to $[a][b] \in K_2^{\mathrm{MW}}(k)$.

Furthermore, there is a natural isomorphism $k^\times \wedge k^\times \cong H_2(k^\times, \mathbb{Z})$ (sending $a \wedge b$ to the homology class $([a]b - [b]a) \otimes 1$) and, by the calculations of Mazzoleni ([11]), these isomorphisms fit into a commutative diagram

$$\begin{array}{ccc} k^\times \wedge k^\times & \xrightarrow{\tilde{\iota}} & K_2^{\mathrm{MW}}(k) \\ \downarrow \cong & & \downarrow \cong \\ H_2(k^\times, \mathbb{Z}) & \xrightarrow{\iota} & H_2(\mathrm{SL}_2(k), \mathbb{Z}) \end{array}$$

where $\tilde{\iota}(a \wedge b) = [a][b] - [b][a] = [a^2][b] = [a][b^2]$.

Furthermore, Mazzoleni has shown that the image of $\tilde{\iota}$ is isomorphic to $2 \cdot K_2^{\mathrm{M}}(k) \subset K_2^{\mathrm{MW}}(k)$ (via

$$\tilde{\iota}(a \wedge b) = [a^2][b] \leftrightarrow \{a^2\}\{b\} = 2\{a\}\{b\}).$$

It follows that the cokernel of $\tilde{\iota}$ is isomorphic to $I^2(k)$, the isomorphism being induced by the homomorphism

$$K_2^{\mathrm{MW}}(k) \rightarrow I^2(k), \quad [a][b] \mapsto \langle\langle a \rangle\rangle \langle\langle b \rangle\rangle.$$

Putting all of this together, we have a natural short exact sequence

$$0 \longrightarrow I^2(k) \longrightarrow \frac{H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z})}{H_2(\mathrm{SL}_2(k), \mathbb{Z})} \xrightarrow{\delta} k^\times \longrightarrow 0.$$

Morel ([14, Theorem 2.15]) shows that if F is a field with discrete valuation v , residue field k and corresponding uniformizer π , there is a well-defined residue homomorphism (depending on the choice of uniformizer) $\delta_\pi^v = \delta_\pi : K_2^{\mathrm{MW}}(F) \rightarrow K_1^{\mathrm{MW}}(k)$ with the properties:

- (1) $\delta_\pi[a][\pi] = [\bar{a}]$ whenever $v(a) = 0$, and
- (2) $\delta_\pi[a][b] = 0$ if $v(a) = v(b) = 0$.

Now let $\Delta : H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \rightarrow K_1^{\mathrm{MW}}(k)$ denote the composite

$$H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \longrightarrow H_2(\mathrm{SL}_2(k(t)), \mathbb{Z}) \xrightarrow{\cong} K_2^{\mathrm{MW}}(k(t)) \xrightarrow{\delta_t} K_1^{\mathrm{MW}}(k).$$

From property (2) of δ_π (with $\pi = t$) it follows that the composite map

$$K_2^{\mathrm{MW}}(k) \longrightarrow K_2^{\mathrm{MW}}(k(t)) \xrightarrow{\delta_t} K_1^{\mathrm{MW}}(k)$$

is the zero map. Thus $\Delta(j(z)) = 0$ for all $z \in H_2(\mathrm{SL}_2(k), \mathbb{Z})$ and hence Δ induces a well-defined homomorphism

$$\tilde{\Delta} : \frac{H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z})}{H_2(\mathrm{SL}_2(k), \mathbb{Z})} \rightarrow K_1^{\mathrm{MW}}(k).$$

To complete the proof of Theorem 4.1 we show that $\tilde{\Delta}$ is an isomorphism.

From our work above, there is a natural injective map $I^2(k) \rightarrow H_2(\mathrm{SL}_2(k), \mathbb{Z})_{-1}$ given by $\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle \mapsto \langle t \rangle j([a, b])$.

On the other hand, there is also a natural short exact sequence

$$0 \longrightarrow I^2(k) \longrightarrow K_1^{\mathrm{MW}}(k) \longrightarrow k^\times \longrightarrow 0$$

where the inclusion $I^2(k) \rightarrow K_1^{\mathrm{MW}}(k)$ is given by $\langle\langle a \rangle\rangle \langle\langle b \rangle\rangle \mapsto \eta[a][b] = [ab] - [a] - [b]$ by (3) above.

Now, from the definition of $\tilde{\Delta}$ and the properties of δ_t , we have

$$\begin{aligned} \tilde{\Delta}(\langle t \rangle j([a, b])) &= \delta_t(\langle t \rangle [a][b]) \\ &= \delta_t(\eta[t][a][b] + [a][b]) \\ &= \delta_t(\eta[a][b][t]) \\ &= \eta[a][b]. \end{aligned}$$

We deduce immediately that there is a natural map of short exact sequences (defining the map $\bar{\Delta}$)

$$\begin{array}{ccccccc} 0 & \longrightarrow & I^2(k) & \longrightarrow & \frac{H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z})}{H_2(\mathrm{SL}_2(k), \mathbb{Z})} & \longrightarrow & k^\times \longrightarrow 0 \\ & & \downarrow = & & \downarrow \tilde{\Delta} & & \downarrow \bar{\Delta} \\ 0 & \longrightarrow & I^2(k) & \longrightarrow & K_1^{\mathrm{MW}}(k) & \longrightarrow & k^\times \longrightarrow 0. \end{array}$$

We now complete the argument by showing that $\bar{\Delta} = \bar{\Delta}_k$ is the identity map on $k^\times$.

We begin by observing that it is enough to show that $\bar{\Delta}_k(a) = a$ for all $a \in (k^\times)^2$ (and for all fields k), since, for any $a \in k^\times$, the diagram

$$\begin{array}{ccc} k^\times & \longrightarrow & k(\sqrt{a})^\times \\ \downarrow \bar{\Delta}_k & & \downarrow \bar{\Delta}_{k(\sqrt{a})} \\ k^\times & \longrightarrow & k(\sqrt{a})^\times \end{array}$$

commutes, and hence $\bar{\Delta}_k(a) = \bar{\Delta}_{k(\sqrt{a})}(a) = a \in k^\times \subset k(\sqrt{a})^\times$.

Recall that δ denotes the connecting homomorphism $H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \rightarrow H_1(k, \mathbb{Z}) = k^\times$ of the Mayer-Vietoris sequence.

For a ring A , we let τ_A denote the composite map

$$A^\times \wedge A^\times \xrightarrow{\cong} H_2(A^\times, \mathbb{Z}) \xrightarrow{\iota} H_2(\mathrm{SL}_2(A), \mathbb{Z}).$$

Proposition 4.2. *Let $A = k[t, t^{-1}]$ and let $a \in k^\times$. Then $\delta(\tau_A(t \wedge a)) = a^2$ in $k^\times$.*

Proof. Let $B_\bullet(G)$ denote the (right) bar resolution of the group $G = \mathrm{SL}_2(k[t, t^{-1}])$. The Mayer-Vietoris sequence is derived from the long exact homology sequence of the exact sequence of complexes

$$0 \rightarrow B_\bullet(G) \otimes_{\mathbb{Z}[G]} \mathbb{Z}[G/\Gamma] \rightarrow B_\bullet(G) \otimes_{\mathbb{Z}[G]} (\mathbb{Z}[G/G_1] \oplus \mathbb{Z}[G/G_2]) \rightarrow B_\bullet(G) \otimes_{\mathbb{Z}[G]} \mathbb{Z} \rightarrow 0.$$

The homology class $\tau_A(t \wedge a) \in H_2(G, \mathbb{Z})$ is represented by the cycle

$$([\iota(t)|\iota(a)] - [\iota(a)|\iota(t)]) \otimes 1 \in B_2(G) \otimes_{\mathbb{Z}[G]} \mathbb{Z}.$$

This lifts to

$$([\iota(a)|\iota(t)] - [\iota(t)|\iota(a)]) \otimes (1 \cdot G_1, 0) \in B_2(G) \otimes_{\mathbb{Z}[G]} (\mathbb{Z}[G/G_1] \oplus \mathbb{Z}[G/G_2]).$$

The boundary homomorphism sends this element to

$$[\iota(a)] \otimes (\iota(t) \cdot G_1 - 1 \cdot G_1, 0) \in B_1(G) \otimes_{\mathbb{Z}[G]} (\mathbb{Z}[G/G_1] \oplus \mathbb{Z}[G/G_2]).$$

Now let

$$w = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \in G_1 = \mathrm{SL}_2(k[t]).$$

Observe that

$$w \cdot \iota(t) = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \cdot \begin{bmatrix} t & 0 \\ 0 & t^{-1} \end{bmatrix} = \begin{bmatrix} 0 & -t^{-1} \\ t & 0 \end{bmatrix} := F(t) \in G_2$$

and hence $\iota(t) = w^{-1}F(t)$ with $w^{-1} \in G_1$, $F(t) \in G_2$.

It follows that $(\iota(t) \cdot G_1 - 1 \cdot G_1, 0)$ is the image of $w^{-1} \cdot (F(t)\Gamma - \Gamma) \in \mathbb{Z}[G/\Gamma]$ under the map $\mathbb{Z}[G/\Gamma] \rightarrow \mathbb{Z}[G/G_1] \oplus \mathbb{Z}[G/G_2]$.

Thus $\delta(\tau_A(a \wedge t))$ is represented by the cycle

$$[\iota(a)] \otimes w^{-1} \cdot (F(t)\Gamma - \Gamma) = ([\iota(a)]\iota(t) - [\iota(a)]w^{-1}) \otimes \Gamma \in B_1(G) \otimes_{\mathbb{Z}[G]} \mathbb{Z}[G/\Gamma].$$

This, in turn, is the image of

$$([\iota(a)]\iota(t) - [\iota(a)]w^{-1}) \otimes 1 \in B_1(G) \otimes_{\mathbb{Z}[\Gamma]} \mathbb{Z}$$

under the natural homology isomorphism $B_\bullet(G) \otimes_{\mathbb{Z}[\Gamma]} \mathbb{Z} \rightarrow B_\bullet(G) \otimes_{\mathbb{Z}[G]} \mathbb{Z}[G/\Gamma]$.

For a group H , we let $C_\bullet(H)$ denote the (right) homogeneous resolution of H . Then the isomorphism $B_\bullet(H) \rightarrow C_\bullet(H)$ is obtained by the map of right $\mathbb{Z}[H]$ -complexes

$$[h_n] \cdots [h_1] \mapsto (h_n \cdot h_{n-1} \cdots h_1, h_{n-1} \cdots h_1, \dots, h_1, 1).$$

Thus $([\iota(a)]\iota(t) - [\iota(a)]w^{-1}) \otimes 1 \in B_1(G) \otimes_{\mathbb{Z}[\Gamma]} \mathbb{Z}$ corresponds to

$$((\iota(at), \iota(t)) - (\iota(a)w^{-1}, w^{-1})) \otimes 1 \in C_1(G) \otimes_{\mathbb{Z}[\Gamma]} \mathbb{Z}.$$

To construct an augmentation preserving map of $\mathbb{Z}[\Gamma]$ -resolutions of $\mathbb{Z}$ from $C_\bullet(G)$ to $C_\bullet(\Gamma)$ we choose a set-theoretic section $s : G/\Gamma \rightarrow G$ and send $(g_n, \dots, g_0)$ to $(\overline{g_n}, \dots, \overline{g_0})$, where $\overline{g} := s(g\Gamma)^{-1}g \in \Gamma$.

To be more specific, we choose a section s satisfying $s(\iota(a)w^{-1}\Gamma) = w^{-1}$ and $s(\iota(at)\Gamma) = \iota(t)$ for all $a \in k^\times$.

Thus the element $(\iota(at), \iota(t)) - (\iota(a)w^{-1}, w^{-1}) \in C_1(G)$ maps to $(\iota(a), 1) - (\iota(a^{-1}), 1) \in C_1(\Gamma)$ (since $w\iota(a)w^{-1} = \iota(a^{-1})$ in G).

Finally the homology class

$$((\iota(a), 1) - (\iota(a^{-1}), 1)) \otimes 1 \in C_1(\Gamma) \otimes_{\mathbb{Z}[\Gamma]} \mathbb{Z}$$

corresponds to $\iota(a) \cdot \iota(a^{-1})^{-1} = \iota(a^2)$ under the isomorphism $H_1(\Gamma, \mathbb{Z}) \cong \Gamma^{\text{ab}}$ and hence to $a^2 \in k^\times \cong \Gamma^{\text{ab}}$. $\square$

Corollary 4.3. *For any field k , we have $\bar{\Delta}_k(a^2) = a^2$ for all $a \in k^\times$.*

Proof. By Proposition 4.2, $a^2 \in k^\times$ is the image, under δ , of the element of $H_2(\text{SL}_2(k), \mathbb{Z})_{-1}$ represented by $\tau_A(a \wedge t)$. Therefore, it is enough to verify that $\Delta(\tau_A(a \wedge t)) = [a^2] \in K_1^{\text{MW}}(k)$:

The image of $\tau_A(a \wedge t)$ in $K_2^{\text{MW}}(k(t))$ is $\tilde{t}(a \wedge t) = [a^2][t]$.

Hence, from the definitions,

$$\bar{\Delta}_k(a^2) = \Delta(\tau_A(a \wedge t)) = \delta_t([a^2][t]) = [a^2]$$

as required. $\square$

5. SOME EXAMPLES AND SPECIAL CASES

The main result of section 4 is that if k is an infinite field of characteristic not equal to 2, then

$$H_2(\text{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \cong H_2(\text{SL}_2(k), \mathbb{Z}) \oplus K_1^{\text{MW}}(k) \cong K_2^{\text{MW}}(k) \oplus K_1^{\text{MW}}(k).$$

This allows us to calculate these homology groups for certain families of fields for which the Milnor K -theory and Witt rings are known.

For a global field F , we will let $\Omega = \Omega(F)$ be the set of real embeddings of F . We denote by $K_2(F)_+$ the kernel of the (split) surjection $K_2(F) \rightarrow \mu_2^\Omega$ induced by the Hilbert symbols associated to each of the real embeddings.

Proposition 5.1. *Let F be a global field. Then there is a natural split exact sequence*

$$0 \rightarrow K_2(F)_+ \rightarrow K_2^{\text{MW}}(F) \rightarrow \mathbb{Z}^\Omega \rightarrow 0.$$

Proof. We begin by recalling from quadratic form theory that for all $n \geq 1$, $I^n(\mathbb{R}) \cong \mathbb{Z}$ with generator $\langle\langle -1 \rangle\rangle \cdots \langle\langle -1 \rangle\rangle$.

For any real embedding $\sigma : F \rightarrow \mathbb{R}$, let T_σ denote the composite homomorphism

$$K_2^{\text{MW}}(F) \rightarrow K_2^{\text{MW}}(\mathbb{R}) \rightarrow I^2(\mathbb{R}) \cong \mathbb{Z}.$$

Thus $T_\sigma([a][b]) = \langle\langle \text{sgn}(\sigma(a)) \rangle\rangle \langle\langle \text{sgn}(\sigma(b)) \rangle\rangle$ where

$$\text{sgn}(x) = \begin{cases} 1, & x > 0 \\ -1, & x < 0. \end{cases}$$

Let

$$T = \oplus_{\sigma \in \Omega} T_\sigma : K_2^{\text{MW}}(F) \rightarrow I^2(\mathbb{R})^\Omega$$

and let $K_2^{\text{MW}}(F)_+ := \text{Ker}(T)$.

Thus we obtain a commutative diagram with exact rows and columns

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & I^3(F) & \xrightarrow{T} & I^3(\mathbb{R})^\Omega & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & K_2^{\text{MW}}(F)_+ & \longrightarrow & K_2^{\text{MW}}(F) & \xrightarrow{T} & I^2(\mathbb{R})^\Omega \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & K_2(F)_+ & \longrightarrow & K_2(F) & \longrightarrow & \mu_2^\Omega \longrightarrow 0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array}$$

where $I^2(\mathbb{R}) \rightarrow \mu_2$ is the unique homomorphism sending $\langle\langle -1 \rangle\rangle$ to -1 .

Now quadratic form theory tells us that, for global fields F , the map $T : I^3(F) \rightarrow I^3(\mathbb{R})^\Omega$ is an isomorphism. The result follows immediately. $\square$

Since $K_2(\mathbb{Q})_+ \cong \bigoplus_{p \text{ odd}} \mathbb{F}_p^\times$ we immediately deduce:

Corollary 5.2. $H_2(\text{SL}_2(\mathbb{Q}), \mathbb{Z}) \cong K_2^{\text{MW}}(\mathbb{Q}) \cong \mathbb{Z} \oplus \left(\bigoplus_{p \text{ odd}} \mathbb{F}_p^\times \right)$.

The situation for $K_1^{\text{MW}}(F)$ is slightly more complicated:

Again, we set

$$K_1^{\text{MW}}(F)_+ = \text{Ker}(K_1^{\text{MW}}(F) \rightarrow I(\mathbb{R})^\Omega)$$

and

$$K_1(F)_+ = F_+^\times = \text{Ker}(F^\times \rightarrow \mu_2^\Omega)$$

(the map being induced by the sgn homomorphisms).

Let $k_2(F) := K_2(F) \otimes \mathbb{Z}/2$ and let $k_2(F)_+ := \text{Ker}(k_2(F) \rightarrow \mu_2^\Omega)$.

Proposition 5.3. *Let F be a global field. There is a split short exact sequences*

$$0 \rightarrow K_1^{\text{MW}}(F)_+ \rightarrow K_1^{\text{MW}}(F) \rightarrow I(\mathbb{R})^\Omega \rightarrow 0$$

and a short exact sequence

$$0 \rightarrow k_2(F)_+ \rightarrow K_1^{\text{MW}}(F)_+ \rightarrow F_+^\times \rightarrow 0$$

which is split if $\Omega \neq \emptyset$.

In particular, if F is a number field admitting a real embedding, as an additive group,

$$K_1^{\text{MW}}(F) \cong \mathbb{Z}^\Omega \oplus k_2(F)_+ \oplus F_+^\times.$$

Proof. Let $\mathcal{T} := \text{Ker}(I^2(F) \rightarrow I^2(\mathbb{R})^\Omega)$. Then there is a commutative diagram with exact rows and columns

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
 & & I^3(F) & \xrightarrow[\cong]{T} & I^3(\mathbb{R})^\Omega & & \\
 & & \downarrow & & \downarrow & & \\
 0 & \longrightarrow & \mathcal{T} & \longrightarrow & I^2(F) & \longrightarrow & I^2(\mathbb{R})^\Omega \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & k_2(F)_+ & \longrightarrow & k_2(F) & \longrightarrow & \mu_2^\Omega \longrightarrow 0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array}$$

It follows that $\mathcal{T} \cong k_2(F)_+$.

Therefore, we obtain a commutative diagram with exact rows and columns

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & k_2(F)_+ & \longrightarrow & I^2(F) & \longrightarrow & I^2(\mathbb{R})^\Omega \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & K_1^{\text{MW}}(F)_+ & \longrightarrow & K_1^{\text{MW}}(F) & \longrightarrow & I(\mathbb{R})^\Omega \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & F_+^\times & \longrightarrow & K^\times & \longrightarrow & \mu_2^\Omega \longrightarrow 0 \\
 & & & & \downarrow & & \downarrow \\
 & & & & 0 & & 0
 \end{array}$$

Finally, if $\Omega \neq \emptyset$, then $F_+^\times$ is a free abelian group and hence the sequence

$$0 \rightarrow k_2(F)_+ \rightarrow K_1^{\text{MW}}(F)_+ \rightarrow F_+^\times \rightarrow 0$$

is split. □

Since $\mathbb{Q}_+^\times \cong \bigoplus_p \mathbb{Z}$ and $k_2(\mathbb{Q})_+ \cong \bigoplus_{p \text{ odd}} \mathbb{Z}/2$ we deduce:

Corollary 5.4. $K_1^{\text{MW}}(\mathbb{Q}) \cong \mathbb{Z} \oplus \left(\bigoplus_p \mathbb{Z} \right) \oplus \left(\bigoplus_{p \text{ odd}} \mathbb{Z}/2 \right)$.

Putting this together with the result of section 4 gives:

Theorem 5.5. $H_2(\text{SL}_2(\mathbb{Q}[t, t^{-1}]), \mathbb{Z}) \cong \mathbb{Z} \oplus \left(\bigoplus_p \mathbb{Z} \right) \oplus \left(\bigoplus_{p \text{ odd}} (\mathbb{F}_p^\times \oplus \mathbb{Z}/2) \right)$.

The results of section 4 also tell us about the stabilization homomorphism from $H_2(\text{SL}_2(k[t, t^{-1}]), \mathbb{Z})$ to $H_2(\text{SL}(k[t, t^{-1}]), \mathbb{Z})$:

Proposition 5.6. *Let k be an infinite field of characteristic not equal to 2. Then the natural stabilization homomorphism*

$$H_2(\text{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \rightarrow H_2(\text{SL}(k[t, t^{-1}]), \mathbb{Z}) = K_2(k[t, t^{-1}])$$

is surjective with kernel isomorphic to $I^3(k) \oplus I^2(k)$.

Proof. By the Fundamental Theorem of Algebraic K -theory (see [20, V.6]) there is a natural split exact sequence

$$0 \rightarrow K_2(k) \rightarrow K_2(k[t, t^{-1}]) \rightarrow K_1(k) \rightarrow 0.$$

Let $\mathcal{K}$ denote the kernel of the map

$$H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \rightarrow H_2(\mathrm{SL}(k[t, t^{-1}]), \mathbb{Z}).$$

Thus we get a natural commutative diagram whose rows are split exact (with compatible splitting maps):

$$\begin{array}{ccccccc}
 & & 0 & & 0 & & 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & I^3(k) & \longrightarrow & \mathcal{K} & \longrightarrow & I^2(k) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & K_2^{\mathrm{MW}}(k) & \longrightarrow & H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) & \longrightarrow & K_1^{\mathrm{MW}}(k) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 0 & \longrightarrow & K_2(k) & \longrightarrow & H_2(\mathrm{SL}(k[t, t^{-1}]), \mathbb{Z}) & \longrightarrow & K_1(k) \longrightarrow 0 \\
 & & \downarrow & & \downarrow & & \downarrow \\
 & & 0 & & 0 & & 0
 \end{array}$$

□

Corollary 5.7. (1) Let k be a quadratically closed field of characteristic not equal to 2. Then the natural stabilization homomorphism

$$H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \rightarrow H_2(\mathrm{SL}(k[t, t^{-1}]), \mathbb{Z})$$

is an isomorphism.

(2) Let k be an infinite field of characteristic not equal to 2 such that $I^3(k) = 0$. Then there is a natural short exact sequence

$$0 \rightarrow k_2(k) \rightarrow H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \rightarrow H_2(\mathrm{SL}(k[t, t^{-1}]), \mathbb{Z}) \rightarrow 0.$$

Proof. (1) When k is quadratically closed $I^n(k) = 0$ for all $n \geq 1$.

(2) In this case

$$I^2(k) = I^2(k)/I^3(k) \cong k_2(k)$$

by Merkurjev's theorem.

□

Remark 5.8. The condition $I^3(k) = 0$ is satisfied, for example, global fields of positive characteristic and by totally imaginary number fields k (and for these fields $K_2(k) \otimes \mathbb{Z}/2\mathbb{Z}$ is infinite). More generally, this condition is satisfied by fields of cohomological dimension 2 or less.

6. SCISSORS CONGRUENCE GROUPS

6.1. The classical scissors congruence group. Let k be a field with at least four elements. The *scissors congruence group*, $\mathcal{P}(k)$, of k is the $\mathbb{Z}$ -module with generators $[a]$, $a \in k^\times$, subject to the relations $[1] = 0$ and

$$[x] - [y] + \left[\frac{y}{x} \right] - \left[\frac{1-x^{-1}}{1-y^{-1}} \right] + \left[\frac{1-x}{1-y} \right], \quad x, y \neq 1.$$

Remark 6.1. When $k = \mathbb{C}$, the minus-eigenspace of $\mathcal{P}(\mathbb{C})$ for the action of complex conjugation can be naturally identified with the scissors congruence group of polyhedra in 3-dimensional hyperbolic space ([2]). This is the origin of the name.

For finite fields we have the following calculation ([5, Lemma 7.4]):

Theorem 6.2. *Let n' denote the odd part of the integer n . Let $\mathbb{F}_q$ denote the finite field with q elements. Then*

$$\mathcal{P}(\mathbb{F}_q) \left[\frac{1}{2} \right] \cong \mathbb{Z}/(q+1)'.$$

We let

$$S_{\mathbb{Z}}^2(k^\times) := \frac{k^\times \otimes_{\mathbb{Z}} k^\times}{\langle \{x \otimes y + y \otimes x \mid x, y \in k^\times\} \rangle},$$

the second (graded) symmetric power. We let $x \circ y$ denote the image of $x \otimes y$ in $S_{\mathbb{Z}}^2(k^\times)$.

The *Bloch group*, $\mathcal{B}(k)$, of the field k is the kernel of the map

$$\lambda : \mathcal{P}(k) \rightarrow S_{\mathbb{Z}}^2(k^\times), \quad [a] \mapsto a \circ (1-a).$$

By a result of Suslin ([19]), the Bloch group of a field is very closely related to the indecomposable K_3 of the field:

Theorem 6.3. *For any field k there is a natural short exact sequence*

$$0 \rightarrow \widetilde{\text{tor}(\mu_k, \mu_k)} \rightarrow K_3^{\text{ind}}(k) \rightarrow \mathcal{B}(k) \rightarrow 0.$$

Here $\widetilde{\text{tor}(\mu_k, \mu_k)} = \text{tor}(\mu_k, \mu_k)$ when $\text{char}(k) = 2$ and otherwise is the nontrivial extension of $\text{tor}(\mu_k, \mu_k)$ by $\mathbb{Z}/2$.

6.2. The refined scissors congruence group. The *refined scissors congruence group*, $\mathcal{RP}(k)$, of k is the $\mathbb{Z}[k^\times/(k^\times)^2]$ -module with generators $[a]$, $a \in k^\times$, subject to the relations $[1] = 0$ and

$$[x] - [y] + \langle x \rangle \left[\frac{y}{x} \right] - \langle x^{-1} - 1 \rangle \left[\frac{1-x^{-1}}{1-y^{-1}} \right] + \langle 1-x \rangle \left[\frac{1-x}{1-y} \right], \quad x, y \neq 1.$$

We endow $S_{\mathbb{Z}}^2(k^\times)$ with the trivial $\mathbb{Z}[k^\times/(k^\times)^2]$ -module structure.

As in [5], we let $\lambda_1 : \mathcal{RP}(k) \rightarrow \mathbb{Z}[k^\times/(k^\times)^2]$ be the $\mathbb{Z}[k^\times/(k^\times)^2]$ -module homomorphism sending $[a]$ to $\langle a \rangle \langle 1-a \rangle$ and let $\lambda_2 : \mathcal{RP}(k) \rightarrow S_{\mathbb{Z}}^2(k^\times)$ be the $\mathbb{Z}[k^\times/(k^\times)^2]$ -homomorphism sending $[a]$ to $a \circ (1-a)$.

Furthermore, let $\Lambda := (\lambda_1, \lambda_2) : \mathcal{RP}(k) \rightarrow \mathbb{Z}[k^\times/(k^\times)^2] \oplus S_{\mathbb{Z}}^2(k^\times)$.

The *refined Bloch group* of k , $\mathcal{RB}(k)$, is the kernel of Λ .

It has the same relation to $H_3(\text{SL}_2(k), \mathbb{Z})$ as $\mathcal{B}(k)$ does to $K_3^{\text{ind}}(k)$:

Theorem 6.4. ([5, Theorem 4.3]) *Let k be a field with at least 28 elements. There is a natural complex*

$$0 \rightarrow \text{tor}(\mu_k, \mu_k) \rightarrow H_3(\text{SL}_2(k), \mathbb{Z}) \rightarrow \mathcal{RB}(k) \rightarrow 0$$

which is exact except possibly at the middle term where the homology is annihilated by 4.

The natural map $\mathcal{RP}(k) \rightarrow \mathcal{P}(k)$ induces a homomorphism $\mathcal{RB}(k) \rightarrow \mathcal{B}(k)$. We denote the kernel of this map by $\mathcal{RB}_0(k)$.

Here we collect some of the relevant facts about $\mathcal{RB}_0(k)$:

Theorem 6.5. *Let k be a field with at least 4 elements.*

- (1) *The map $\mathcal{RB}(k) \rightarrow \mathcal{B}(k)$ is surjective. Hence there is a natural short exact sequence of $\mathbb{Z}[k^\times/(k^\times)^2]$ -modules*

$$0 \rightarrow \mathcal{RB}_0(k) \rightarrow \mathcal{RB}(k) \rightarrow \mathcal{B}(k) \rightarrow 0.$$

Furthermore, if k has finitely many square classes, after tensoring with $\mathbb{Z}[\frac{1}{2}]$ this sequence is split.

- (2) *If k has at least 28 elements there is a natural short exact sequence of $\mathbb{Z}[k^\times/(k^\times)^2]$ -modules*

$$0 \rightarrow \mathcal{RB}_0(k) \left[\frac{1}{2} \right] \rightarrow H_3(\mathrm{SL}_2(k), \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow K_3^{\mathrm{ind}}(k) \left[\frac{1}{2} \right] \rightarrow 0.$$

Furthermore, if k has finitely many square classes, this sequence is split.

- (3) *For any infinite field k there is a natural exact sequence*

$$0 \rightarrow \mathcal{RB}_0(k) \left[\frac{1}{2} \right] \rightarrow H_3(\mathrm{SL}_2(k), \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow H_3(\mathrm{SL}_n(k), \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow K_3^{\mathrm{M}}(k) \left[\frac{1}{2} \right] \rightarrow 0$$

for any $n \geq 3$.

- (4) *If k is finite or real-closed or quadratically closed, then $\mathcal{RB}_0(k) = 0$.*
 (5) *Suppose that k is a local field with finite residue field $\bar{k}$ of odd order. If $\mathbb{Q}_3 \subset k$ we suppose that $[k : \mathbb{Q}_3]$ is odd. Then there is a natural isomorphism*

$$\mathcal{RB}_0(k) \left[\frac{1}{2} \right] \cong \mathcal{P}(\bar{k}) \left[\frac{1}{2} \right].$$

- (6) *Suppose that k is the field of fractions of a unique factorization domain A . Let $\mathcal{P}$ be a set of representatives of the association classes of prime elements of $\mathcal{P}$. Then there is a natural surjective homomorphism*

$$\mathcal{RB}_0(k) \left[\frac{1}{2} \right] \twoheadrightarrow \bigoplus_{p \in \mathcal{P}} \mathcal{P}(\bar{k}_p) \left[\frac{1}{2} \right].$$

Proof. (1) This is [5, Corollary 5.1].

- (2) This is [5, Lemma 5.2].

The statement about the splitting of the sequences follows from the fact that $\mathcal{RB}_0(k) \left[\frac{1}{2} \right] = \mathcal{I}_k \mathcal{RB}(k) \left[\frac{1}{2} \right]$, together with the fact that if $k^\times/(k^\times)^2$ is finite and if M is any $\mathbb{Z}[k^\times/(k^\times)^2] \left[\frac{1}{2} \right]$ -module, then the sequence

$$0 \rightarrow \mathcal{I}_k M \rightarrow M \rightarrow M/\mathcal{I}_k M \rightarrow 0$$

is naturally split.

- (3) This follows from [6, Theorem 4.7] and [5, Lemma 5.2].
 (4) For quadratically closed field the is immediate from the definition. For real closed fields this is the result of Parry and Sah [15]. For finite fields, this is [5, Lemma 7.1].
 (5) This is [4, Theorem 6.19].
 (6) This is [4, Theorem 5.1].

□

6.3. **The module $\mathcal{RP}_1(k)$.** The $\mathbb{Z}[k^\times/(k^\times)^2]$ -module

$$\mathcal{RP}_1(k) := \text{Ker}(\lambda_1 : \mathcal{RP}(k) \rightarrow S_{\mathbb{Z}}^2(k^\times))$$

will play a key role in our calculations below.

Note that $\mathcal{RB}(k)$ is the kernel of $\lambda_2 : \mathcal{RP}_1(k) \rightarrow S_{\mathbb{Z}}^2(k^\times)$.

Lemma 6.6. *There is a natural short exact sequence of $\mathbb{Z}[k^\times/(k^\times)^2]$ -modules*

$$0 \rightarrow \mathcal{RB}_0(k) \left[\frac{1}{2} \right] \rightarrow \mathcal{RP}_1(k) \left[\frac{1}{2} \right] \rightarrow \mathcal{P}(k) \left[\frac{1}{2} \right] \rightarrow 0$$

which is split if k has finitely many square classes.

Proof. For $x \in k^\times$ we let $\psi_1(x)$ denote the element $[x] + \langle -1 \rangle [x^{-1}] \in \mathcal{RP}(k)$.

An elementary calculation (see, for example, [4, Lemma 3.3]) shows that for any $1 \neq x \in k^\times$

$$r(x) := (\langle -1 \rangle + 1)[x] + \langle\langle 1 - x \rangle\rangle \psi_1(x) \in \text{Ker}(\lambda_1) = \mathcal{RP}_1(k).$$

But $\lambda_2(r(x)) = 2\lambda([x])$.

Let

$$\mathcal{R}(k) := \text{Im}(\lambda : \mathcal{P}(k) \rightarrow S_{\mathbb{Z}}^2(k^\times)) \text{ and } \mathcal{R}_1(k) := \text{Im}(\lambda_2 : \mathcal{RP}_1(k) \rightarrow S_{\mathbb{Z}}^2(k^\times)).$$

Then $\mathcal{R}_1(k) \subset \mathcal{R}(k)$ and the quotient, $C(k)$, is annihilated by 2.

The statement now follows by applying the snake lemma to the map of short exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{RB}(k) & \longrightarrow & \mathcal{RP}_1(k) & \longrightarrow & \mathcal{R}_1(k) \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{B}(k) & \longrightarrow & \mathcal{P}(k) & \longrightarrow & \mathcal{R}(k) \longrightarrow 0 \end{array}$$

and tensoring with $\mathbb{Z} \left[\frac{1}{2} \right]$. □

We let $\mathcal{K}_k^{(1)}$ denote the $\mathbb{Z}[k^\times/(k^\times)^2]$ -submodule of $\mathcal{RP}(k)$ generated by the elements $\psi_1(x)$, $x \in k^\times$.

Let $\widetilde{\mathcal{RP}}(k)$ denote the $\mathbb{Z}[k^\times/(k^\times)^2]$ -module $\mathcal{RP}(k)/\mathcal{K}_k^{(1)}$. Let $\widetilde{\mathcal{RP}}_1(k)$ denote the image of the map $\mathcal{RP}_1(k) \rightarrow \widetilde{\mathcal{RP}}(k)$.

In [4, Lemma 3.3] it is shown that $\mathcal{RP}_1(k) \cap \mathcal{K}_k^{(1)}$ is precisely the torsion subgroup of $\mathcal{K}_k^{(1)}$ and is annihilated by 4. We deduce:

Lemma 6.7. *The natural map $\mathcal{RP}_1(k) \rightarrow \widetilde{\mathcal{RP}}_1(k)$ induces an isomorphism*

$$\mathcal{RP}_1(k) \left[\frac{1}{2} \right] \cong \widetilde{\mathcal{RP}}_1(k) \left[\frac{1}{2} \right].$$

6.4. **Specialization homomorphisms.** Suppose given a field F with valuation $v : F^\times \rightarrow \Gamma$.

Let O_v be the corresponding valuation ring, let U_v the group of units of O_v and let k be the residue field. Given $a \in U_v$, we denote by $\bar{a}$ its image in k .

Given an $\mathbb{Z}[k^\times/(k^\times)^2]$ -module M , we let M_F denote the $\mathbb{Z}[F^\times/(F^\times)^2]$ -module

$$\mathbb{Z}[F^\times/(F^\times)^2] \otimes_{\mathbb{Z}[U_v/U_v^2]} M.$$

The following is [4, Theorem 4.9] :

Theorem 6.8. *There is a natural $\mathbb{Z}[F^\times/(F^\times)^2]$ -module homomorphism $S_v : \mathcal{RP}(F) \rightarrow \widetilde{\mathcal{RP}}(k)_F$ which sends $[a]$ to $1 \otimes [\bar{a}]$ when $a \in U_v$.*

It can be easily verified that S_v restricts to a map $S_v : \widetilde{\mathcal{RP}}_1(F) \rightarrow \widetilde{\mathcal{RP}}_1(k)_F$ (see [3, Section 5.2]).

When v is a discrete value and π is a uniformizer, then there is a $\mathbb{Z}[U_v/U_v^2]$ -module decomposition $\mathbb{Z}[F^\times/(F^\times)^2] \cong \mathbb{Z}[U_v/U_v^2] \oplus \langle \pi \rangle \cdot \mathbb{Z}[U_v/U_v^2]$ where $\langle \pi \rangle \in F^\times/(F^\times)^2$ is the square class of π . Thus for any $\mathbb{Z}[k^\times/(k^\times)^2]$ -module M , there is a $\mathbb{Z}[U_v/U_v^2]$ -decomposition (depending on the choice of π)

$$M_F \cong M \oplus \langle \pi \rangle \cdot M.$$

We let $\rho_\pi : M_F \rightarrow M$ denote the resulting $\mathbb{Z}[U_v/U_v^2]$ -map

$$M_F \rightarrow \langle \pi \rangle \cdot M \cong M$$

arising from projection on the second factor.

When v is a discrete value and π is a uniformizer, we let δ_π denote the composite

$$\mathcal{RP}_1(F) \xrightarrow{S_v} \widetilde{\mathcal{RP}}_1(k)_F \xrightarrow{\rho_\pi} \widetilde{\mathcal{RP}}_1(k).$$

7. THE THIRD HOMOLOGY OF $\mathrm{SL}_2(k[t, t^{-1}])$

In this section we prove the following result:

Theorem 7.1. *Let k be an infinite field. Then there is a natural isomorphism*

$$H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\tfrac{1}{2}]) \cong H_3(\mathrm{SL}_2(k), \mathbb{Z}[\tfrac{1}{2}]) \oplus \mathcal{RP}_1(k)[\tfrac{1}{2}].$$

7.1. Preliminaries. Let G be a group and $L_\bullet$ a nonnegative acyclic complex of $\mathbb{Z}[G]$ -modules augmented over $\mathbb{Z}$ via a map $\epsilon : L_0 \rightarrow \mathbb{Z}$; i.e. ϵ induces a weak equivalence of complexes of $\mathbb{Z}[G]$ -modules $L_\bullet \simeq \mathbb{Z}[0]$.

If we suppose further that each of the $\mathbb{Z}$ -modules L_n is free then for any abelian group M , $\epsilon \otimes \mathrm{Id}_M$ induces a weak equivalence of complexes

$$L_\bullet \otimes_{\mathbb{Z}} M \simeq \mathbb{Z}[0] \otimes_{\mathbb{Z}} M = M[0].$$

If M is furthermore a $\mathbb{Z}[G]$ -module, then this is a weak equivalence of complexes of $\mathbb{Z}[G]$ -modules if G acts diagonally on $L_\bullet \otimes_{\mathbb{Z}} M$. It follows that there is an induced isomorphism of homology groups

$$H_n(G, L_\bullet \otimes_{\mathbb{Z}} M) \cong H_n(G, M).$$

Here the left-hand term is the hyperhomology of G with coefficients in the complex $L_\bullet \otimes_{\mathbb{Z}} M$. This, by definition, is the homology of the total complex $T_\bullet(G, M)$ associated to the double complex $D_{p,q}(G, M) = B_p(G) \otimes_{\mathbb{Z}[G]} (L_q \otimes_{\mathbb{Z}} M)$.

Associated to the double complex $D_{p,q}(G, M)$ is a spectral sequence of the form

$$E_{p,q}^1(G, M) = H_p(G, L_q \otimes_{\mathbb{Z}} M) \Rightarrow H_{p+q}(G, M).$$

We make the following observations about this construction:

- (1) **Functoriality:** Given a map of pairs $(H, M') \rightarrow (G, M)$, and endowing $L_\bullet$ with the structure of a $\mathbb{Z}[H]$ -complex via the map $H \rightarrow G$, we obtain a natural maps of complexes $D_{p,q}(H, M') \rightarrow D_{p,q}(G, M)$ and $T_\bullet(H, M') \rightarrow T_\bullet(G, M)$.
- (2) The functor $M \mapsto L_\bullet \otimes M$ from $\mathbb{Z}[G]$ modules to $\mathbb{Z}[G]$ -complexes is exact since the L_q are $\mathbb{Z}$ -free, and hence so are the functors $M \mapsto \{D_{p,q}(G, M)\}_{p,q}$ and $M \mapsto T_\bullet(G, M)$.
- (3) The natural map $D_{\bullet,0}(G, M) \rightarrow T_\bullet(G, M)$ is a map of complexes for which the resulting edge homomorphism on homology $H_\bullet(G, L_0 \otimes M) \rightarrow H_\bullet(G, M)$ coincides with that induced by $\epsilon \otimes \mathrm{Id}_M$.

(4) Given subgroups $H \subset K \subset G$, there are natural composite maps of double complexes

$$D_{p,q}(H, \mathbb{Z}) \rightarrow D_{p,q}(H, \mathbb{Z}[G/K]) \rightarrow D_{p,q}(G, \mathbb{Z}[G/K]).$$

Here, the first map is induced from the inclusion of $\mathbb{Z}[H]$ -modules

$$\mathbb{Z} \rightarrow \mathbb{Z}[G/K], 1 \mapsto K.$$

Thus it sends an element of the form $z \otimes \ell \in D_{p,q}(H, \mathbb{Z}) = B_p(H) \otimes L_q$ to $z \otimes (\ell \otimes K) \in D_{p,q}(H, \mathbb{Z}[G/K]) = B_p(H) \otimes (L_q \otimes \mathbb{Z}[G/K])$. For convenience, we will use the following notation: If $w \in D_{p,q}(H, \mathbb{Z})$ we will let $w \otimes K$ denote its image in $D_{p,q}(H, \mathbb{Z}[G/K])$ or $D_{p,q}(G, \mathbb{Z}[G/K])$.

In this article, the relevant example is the case where $G = \mathrm{SL}_2(k)$ and L_q is the free abelian group of $(q+1)$ -tuples of distinct elements of $\mathbb{P}^1(k)$ (with the usual action of $\mathrm{SL}_2(k)$ on $\mathbb{P}^1(k)$). The boundary map $d_q : L_{q+1} \rightarrow L_q$ is the standard simplicial boundary map.

We will require the following facts about the associated spectral sequence

$$E_{p,q}^1 := E_{p,q}^1(\mathrm{SL}_2(k), \mathbb{Z}) \Rightarrow H_{p+q}(\mathrm{SL}_2(k), \mathbb{Z}) :$$

Lemma 7.2. *The map*

$$\begin{aligned} B_p(k^\times) \otimes_{\mathbb{Z}[k^\times]} \mathbb{Z} &\rightarrow B_p(\mathrm{SL}_2(k)) \otimes_{\mathbb{Z}[\mathrm{SL}_2(k)]} L_0 = D_{p,0} \\ z \otimes 1 &\mapsto z \otimes (\infty) \end{aligned}$$

induces an isomorphism

$$H_p(k^\times, \mathbb{Z}) \cong E_{p,0}^1 = H_p(\mathrm{SL}_2(k), L_0).$$

With this identification, the edge homomorphism

$$H_p(k^\times, \mathbb{Z}) \cong E_{p,0}^1 \longrightarrow E_{p,0}^\infty \hookrightarrow H_p(\mathrm{SL}_2(k), \mathbb{Z})$$

is identified with the map ι . In particular, $E_{p,0}^\infty \cong H_p(k^\times, \mathbb{Z})/\mathrm{Ker}(\iota)$.

Proof. L_0 is a transitive permutation module over $\mathrm{SL}_2(k)$ and the stabilizer of (∞) is the subgroup $B = B(k)$ of upper-triangular matrices. By Shapiro's Lemma it follows that the map of complexes

$$B_\bullet(B) \otimes_{\mathbb{Z}[B]} \mathbb{Z} \rightarrow D_{\bullet,0}, \quad z \otimes 1 \mapsto z \otimes (\infty)$$

induces an isomorphism on homology $H_\bullet(B, \mathbb{Z}) \cong H_\bullet(\mathrm{SL}_2(k), L_0)$.

However, for any infinite field k , the natural inclusion $k^\times \rightarrow B$ induces an isomorphism on homology $H_\bullet(k^\times, \mathbb{Z}) \cong H_\bullet(B, \mathbb{Z})$ by Theorem 2.3 above. $\square$

Lemma 7.3. (1) $2 \cdot E_{2,1}^\infty = 0 = 2 \cdot E_{1,2}^\infty$ and hence $E_{2,1}^\infty \left[\frac{1}{2} \right] = 0 = E_{1,2}^\infty \left[\frac{1}{2} \right]$

$$(2) \quad E_{0,3}^3 \left[\frac{1}{2} \right] \cong \mathcal{RP}_1(k) \left[\frac{1}{2} \right].$$

$$(3) \quad E_{2,0}^3 \left[\frac{1}{2} \right] \cong H_2(k^\times, \mathbb{Z} \left[\frac{1}{2} \right]) \cong (k^\times \wedge k^\times) \left[\frac{1}{2} \right] \text{ and if we let } \rho \text{ denote the isomorphism}$$

$$(k^\times \wedge k^\times) \left[\frac{1}{2} \right] \rightarrow S_{\mathbb{Z}}^2(k^\times) \left[\frac{1}{2} \right], \quad a \wedge b \mapsto 2(a \circ b)$$

then we have an equality of maps

$$\rho \circ d^3 = \lambda_2 : \mathcal{RP}_1(k) \left[\frac{1}{2} \right] = E_{3,0}^3 \rightarrow S_{\mathbb{Z}}^2(k^\times) \left[\frac{1}{2} \right].$$

The proofs of these facts can be found in section 4 of [5] (where low-dimensional terms of the spectral sequence associated to $D_{p,q}(\mathrm{SL}_2(k), \mathbb{Z})$ are calculated).

In particular, we have the following:

Lemma 7.4. $E_{3,0}^\infty \left[\frac{1}{2} \right] \cong \mathcal{RB}(k) \left[\frac{1}{2} \right]$ and the inclusion $k^\times \rightarrow \mathrm{SL}_2(k)$ gives rise to a natural exact sequence

$$H_3(k^\times, \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow H_3(\mathrm{SL}_2(k), \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow \mathcal{RB}(k) \left[\frac{1}{2} \right] \rightarrow 0.$$

7.2. The Mayer-Vietoris sequence again. When $i = 3$ in the Mayer-Vietoris sequence, we obtain an exact sequence

$$H_3(k^\times, \mathbb{Z}) \longrightarrow H_3(\mathrm{SL}_2(k), \mathbb{Z}) \oplus H_2(\mathrm{SL}_2(k), \mathbb{Z}) \longrightarrow H_2(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \xrightarrow{\delta} H_2(k^\times, \mathbb{Z}) \longrightarrow \dots$$

Thus, tensoring with $\mathbb{Z} \left[\frac{1}{2} \right]$ and using Lemma 7.4 we obtain a natural exact sequence

$$0 \longrightarrow \mathcal{RB}(k) \left[\frac{1}{2} \right] \xrightarrow{\beta} \frac{H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right])}{H_3(\mathrm{SL}_2(k), \mathbb{Z} \left[\frac{1}{2} \right])} \xrightarrow{\delta} H_2(k^\times, \mathbb{Z} \left[\frac{1}{2} \right]) \longrightarrow H_2(\mathrm{SL}_2(F), \mathbb{Z} \left[\frac{1}{2} \right])$$

Here the map β is calculated as follows: Given $x \in \mathcal{RB}(k) \left[\frac{1}{2} \right]$, choose $z \in H_3(\mathrm{SL}_2(k), \mathbb{Z} \left[\frac{1}{2} \right])$ mapping to x . Then $\beta(x)$ is the class of $\langle t \rangle \cdot j(z)$ in

$$\frac{H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right])}{H_3(\mathrm{SL}_2(k), \mathbb{Z} \left[\frac{1}{2} \right])}.$$

7.3. The map $H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow \widetilde{\mathcal{RP}}_1(k) \left[\frac{1}{2} \right]$. Now let $\Delta : H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \rightarrow \widetilde{\mathcal{RP}}_1(k)$ denote the following composition of maps:

$$H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \longrightarrow H_3(\mathrm{SL}_2(k(t)), \mathbb{Z}) \longrightarrow \mathcal{RB}(k(t)) \longrightarrow \widetilde{\mathcal{RP}}_1(k(t)) \xrightarrow{\delta_t} \widetilde{\mathcal{RP}}_1(k).$$

The composite map

$$\mathcal{RP}_1(k) \longrightarrow \mathcal{RP}_1(k(t)) \xrightarrow{S_v} \widetilde{\mathcal{RP}}_1(k)_F = \widetilde{\mathcal{RP}}_1(k) \oplus \langle t \rangle \cdot \widetilde{\mathcal{RP}}_1(k)$$

is just the projection on the first factor (since $S_v([a]) = 1 \otimes [\bar{a}]$ when $a \in U_v$) and hence the composite

$$\mathcal{RP}_1(k) \longrightarrow \mathcal{RP}_1(k(t)) \xrightarrow{\delta_t} \widetilde{\mathcal{RP}}_1(k)$$

is the zero map.

From the commutativity of the diagram

$$\begin{array}{ccccc} & & H_3(\mathrm{SL}_2(k), \mathbb{Z}) & \longrightarrow & \mathcal{RB}(k) \\ & \swarrow j & \downarrow & & \downarrow \\ H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}) & \longrightarrow & H_3(\mathrm{SL}_2(k(t)), \mathbb{Z}) & \longrightarrow & \mathcal{RB}(k(t)) \end{array}$$

it follows that $\Delta(j(z)) = 0$ for all $z \in H_3(\mathrm{SL}_2(k), \mathbb{Z})$ and hence there is an induced map

$$\tilde{\Delta} : \frac{H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z})}{H_3(\mathrm{SL}_2(k), \mathbb{Z})} \rightarrow \widetilde{\mathcal{RP}}_1(k).$$

We will show that, at least after tensoring with $\mathbb{Z} \left[\frac{1}{2} \right]$, $\tilde{\Delta}$ is an isomorphism.

We recall that the map

$$k^\times \wedge k^\times \cong H_2(k^\times, \mathbb{Z}) \xrightarrow{\iota} H_2(\mathrm{SL}_2(k), \mathbb{Z}) \cong K_2^{\mathrm{MW}}(k)$$

factors as

$$k^\times \wedge k^\times \xrightarrow{\tilde{\sigma}} 2 \cdot K_2^M(k) \hookrightarrow K_2^{MW}(k)$$

where $\tilde{\sigma}(a \wedge b) = 2\{a\}\{b\}$. In particular,

$$\text{Ker}(\iota) = \text{Ker}(\tilde{\sigma}) = \text{Im}(\delta : H_3(\text{SL}_2(k[t, t^{-1}]), \mathbb{Z}) \rightarrow H_2(k^\times, \mathbb{Z})).$$

(In fact, this is, of course, the subgroup of $k^\times \wedge k^\times$ generated by terms of the form $(1 - a) \wedge a$, where $a \neq 0, 1$.)

The square

$$\begin{array}{ccc} k^\times \wedge k^\times & \xrightarrow{\tilde{\sigma}} & 2 \cdot K_2^M(k) \\ \rho \downarrow & & \downarrow \\ S_{\mathbb{Z}}^2(k^\times) & \xrightarrow{\sigma} & K_2^M(k) \end{array}$$

– where σ is the symbol map $a \circ b \mapsto \{a\}\{b\}$ – commutes, and the vertical maps become isomorphisms upon tensoring with $\mathbb{Z}[\frac{1}{2}]$.

Thus ρ induces an isomorphism

$$\text{Ker}(\tilde{\sigma}) \left[\frac{1}{2} \right] \cong \text{Ker}(\sigma) \left[\frac{1}{2} \right].$$

We have a diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{RB}(k) \left[\frac{1}{2} \right] & \xrightarrow{\beta} & \frac{H_3(\text{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}])}{H_3(\text{SL}_2(k), \mathbb{Z}[\frac{1}{2}])} & \xrightarrow{\delta} & \text{Ker}(\tilde{\sigma}) \left[\frac{1}{2} \right] \longrightarrow 0 \\ & & \downarrow \text{Id} & & \downarrow \tilde{\Delta} & & \downarrow \rho \\ 0 & \longrightarrow & \mathcal{RB}(k) \left[\frac{1}{2} \right] & \longrightarrow & \mathcal{RP}_1(k) \left[\frac{1}{2} \right] & \xrightarrow{\lambda_2} & \text{Ker}(\sigma) \left[\frac{1}{2} \right] \longrightarrow 0. \end{array}$$

To prove that $\tilde{\Delta}$ is an isomorphism, it is enough to show that both squares in this diagram commute.

Let $x \in \mathcal{RB}(k) \left[\frac{1}{2} \right]$. Let $z \in H_3(\text{SL}_2(k), \mathbb{Z}[\frac{1}{2}])$ map to x . Then $\beta(x)$ is represented by the element $\langle t \rangle \cdot j(z) \in H_3(\text{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}])$. This element maps to $\langle t \rangle \cdot x \in \mathcal{RB}(k(t)) \left[\frac{1}{2} \right]$ under the composite

$$H_3(\text{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \rightarrow H_3(\text{SL}_2(k(t)), \mathbb{Z}[\frac{1}{2}]) \rightarrow \mathcal{RB}(k(t)) \left[\frac{1}{2} \right].$$

This element, in turn, maps to $x \in \mathcal{RB}(k) \left[\frac{1}{2} \right] \subset \mathcal{RB}(k(t)) \left[\frac{1}{2} \right]$ under the map δ_t . This shows that the left-hand square commutes.

Finally, the commutativity of the right-hand square follows from

Lemma 7.5. *For any infinite field k , the diagram*

$$\begin{array}{ccc} H_3(\text{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) & \xrightarrow{\delta} & (k^\times \wedge k^\times) \left[\frac{1}{2} \right] \\ \downarrow \Delta & & \downarrow \rho \\ \mathcal{RP}_1(k) \left[\frac{1}{2} \right] & \xrightarrow{\lambda_2} & S_{\mathbb{Z}}^2(k^\times) \left[\frac{1}{2} \right] \end{array}$$

commutes.

Proof. Recall that the Mayer-Vietoris exact sequence associated to the amalgamated product decomposition

$$G = \mathrm{SL}_2(k[t, t^{-1}]) = \mathrm{SL}_2(k[t]) \star_{\Gamma} \mathrm{SL}_2(k[t])^{A(t)} = G_1 \star_{\Gamma} G_2$$

is the long exact homology sequence associated to the short exact sequence of $\mathbb{Z}[G]$ -modules

$$0 \rightarrow \mathbb{Z}[G/\Gamma] \rightarrow \mathbb{Z}[G/G_1] \oplus \mathbb{Z}[G/G_2] \rightarrow \mathbb{Z} \rightarrow 0.$$

This is therefore the long exact sequence resulting from the short exact sequence of complexes

$$0 \rightarrow T(G, \mathbb{Z}[G/\Gamma]) \rightarrow T(G, \mathbb{Z}[G/G_1]) \oplus T(G, \mathbb{Z}[G/G_2]) \rightarrow T(G, \mathbb{Z}) \rightarrow 0$$

(where G acts on the complex $L_{\bullet}$ via the map $G \rightarrow \mathrm{SL}_2(k)$ obtained by sending t to 1).

Suppose given $w \in \mathrm{Ker}(\iota : H_2(k^{\times}, \mathbb{Z}[\frac{1}{2}]) \rightarrow H_2(G_1, \mathbb{Z}[\frac{1}{2}])) = \mathrm{Ker}(\tilde{\sigma})$. Suppose further that w is represented by the cycle $z \otimes 1 \in B_2(k^{\times}) \otimes_{\mathbb{Z}[k^{\times}]} \mathbb{Z}[\frac{1}{2}]$.

Let $\tilde{z} = (0, 0, z \otimes (\infty))$ be the corresponding element of

$$T_2(k^{\times}, \mathbb{Z}[\frac{1}{2}]) = \bigoplus_{p+q=2} D_{p,q}(k^{\times}, \mathbb{Z}[\frac{1}{2}]) = \bigoplus_{p=0}^2 (B_p(k^{\times}) \otimes_{\mathbb{Z}[k^{\times}]} (L_{2-p}[\frac{1}{2}])).$$

Since w maps to 0 in $H_2(G_1, \mathbb{Z}[\frac{1}{2}])$, it follows that the image of $\tilde{z}$ in $T_2(G_1, \mathbb{Z}[\frac{1}{2}])$ is a boundary. Thus there exists $x = (x_0, x_1, x_2, x_3) \in T_3(G_1, \mathbb{Z}[\frac{1}{2}])$ satisfying $dx = \tilde{z}$.

Since the first two components of $\tilde{z}$ are zeroes, it follows that x_0 represents an element, α , of $E_{0,3}^3 = \mathcal{RP}_1(k)[\frac{1}{2}]$ and that $d^3(\alpha) \in E_{2,0}^3 \cong H_2(k^{\times}, \mathbb{Z}[\frac{1}{2}])$ is represented by $(\mathrm{Id}_{B_2(G_1)} \otimes d)(x_2) \in B_2(G_1) \otimes L_0[\frac{1}{2}]$. However, $(\mathrm{Id}_{B_2} \otimes d)(x_2) \pm (d \otimes \mathrm{Id}_{L_0})(x_3) = z \otimes (\infty)$. Hence, $d^3(\alpha) = w$ in $H_2(k^{\times}, \mathbb{Z}[\frac{1}{2}]) \cong H_2(G_1, L_0[\frac{1}{2}])$.

Let $C_t : G_1 \rightarrow G_2$ denote the isomorphism given by conjugating by $A(t)$: $C_t(g) = A(t)^{-1}gA(t) = g^{A(t)}$. Observe that C_t induces the identity map of the diagonal subgroup $k^{\times}$, since $A(t)$ commutes with other diagonal matrices.

Thus $C_t(x) = (C_t(x_0), C_t(x_1), C_t(x_2), C_t(x_3)) \in T_3(G_2, \mathbb{Z}[\frac{1}{2}])$. Then $C_t(x_0)$ represents an element, $C_t(\alpha)$, of $E_{0,3}^3 = C_t(\mathcal{RP}_1(k))[\frac{1}{2}]$ and $d^3(C_t(\alpha)) = w$ in $E_{2,0}^3 = H_2(k^{\times}, \mathbb{Z}[\frac{1}{2}])$.

Now the cycle $\tilde{z} \in T_2(k^{\times}, \mathbb{Z}[\frac{1}{2}])$ maps to the cycle $\tilde{z} \otimes \Gamma \in T_2(G, \mathbb{Z}[G/\Gamma][\frac{1}{2}])$, and this cycle in turn represents the class $w \in H_2(k^{\times}, \mathbb{Z}[\frac{1}{2}]) \cong H_2(\Gamma, \mathbb{Z}[\frac{1}{2}])$.

Under the map

$$T_2(G, \mathbb{Z}[G/\Gamma][\frac{1}{2}]) \rightarrow T_2(G, \mathbb{Z}[G/G_1][\frac{1}{2}]) \oplus T_2(G, \mathbb{Z}[G/G_2][\frac{1}{2}])$$

$\tilde{z} \otimes \Gamma$ maps to $(\tilde{z} \otimes G_1, \tilde{z} \otimes G_2)$. By the calculations above, this is the boundary of

$$(x \otimes G_1, C_t(x) \otimes G_2) \in T_3(G, \mathbb{Z}[G/G_1][\frac{1}{2}]) \oplus T_3(G, \mathbb{Z}[G/G_2][\frac{1}{2}]).$$

This element in turn maps to the cycle $W = C_t(x) - x \in T_3(G, \mathbb{Z}[\frac{1}{2}])$ under the map

$$T_3(G, \mathbb{Z}[G/G_1][\frac{1}{2}]) \oplus T_3(G, \mathbb{Z}[G/G_2][\frac{1}{2}]) \rightarrow T_3(G, \mathbb{Z}[\frac{1}{2}]).$$

By construction, the cycle W represents an element of $H_3(G, \mathbb{Z}[\frac{1}{2}])$ which maps to $w \in H_2(k^{\times}, \mathbb{Z}[\frac{1}{2}])$ under the connecting homomorphism of the Mayer-Vietoris sequence of the amalgamated product.

Now the image of W in $T_3(\mathrm{SL}_2(k(t)), \mathbb{Z}[\frac{1}{2}])$ represents a homology class in $H_3(\mathrm{SL}_2(k(t)), \mathbb{Z}[\frac{1}{2}])$ which maps to $\langle t \rangle \alpha - \alpha \in \mathcal{RB}(k(t))[\frac{1}{2}]$. By definition, this maps to the class $\alpha \in \widetilde{\mathcal{RP}}_1(k)[\frac{1}{2}]$ under the composite

$$\mathcal{RB}(k(t)) \xrightarrow{S_t} \widetilde{\mathcal{RP}}_1(k)_{k(t)} \xrightarrow{\rho_t} \widetilde{\mathcal{RP}}_1(k).$$

Thus $\Delta(W) = \alpha$ and hence

$$\lambda_2(\Delta(W)) = \lambda_2(\alpha) = \rho(d^3(\alpha)) = \rho(w) = \rho(\delta(W))$$

as required. $\square$

8. SOME EXAMPLES AND SPECIAL CASES

The main result of section 7 is that

$$H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \cong H_3(\mathrm{SL}_2(k), \mathbb{Z}[\frac{1}{2}]) \oplus \mathcal{RP}_1(k)[\frac{1}{2}].$$

By Theorem 6.5 (2) and Lemma 6.6 there are natural short exact sequences

$$0 \rightarrow \mathcal{RB}_0(k)[\frac{1}{2}] \rightarrow H_3(\mathrm{SL}_2(k), \mathbb{Z}[\frac{1}{2}]) \rightarrow K_3^{\mathrm{ind}}(k)[\frac{1}{2}] \rightarrow 0$$

and

$$0 \rightarrow \mathcal{RB}_0(k)[\frac{1}{2}] \rightarrow \mathcal{RP}_1(k)[\frac{1}{2}] \rightarrow \mathcal{P}(k)[\frac{1}{2}] \rightarrow 0.$$

Thus we have:

Proposition 8.1. *Let k be an infinite field. Then there is a natural surjective homomorphism*

$$H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \rightarrow K_3^{\mathrm{ind}}(k)[\frac{1}{2}] \oplus \mathcal{P}(k)[\frac{1}{2}]$$

whose kernel is isomorphism to $\mathcal{RB}_0(k)^{\oplus 2}$.

By Theorem 6.5 (4), we deduce:

Corollary 8.2. *Let k be a quadratically closed or real-closed field. Then there is a natural isomorphism*

$$H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \cong K_3^{\mathrm{ind}}(k)[\frac{1}{2}] \oplus \mathcal{P}(k)[\frac{1}{2}].$$

By Theorem 6.5 (5) we have:

Corollary 8.3. *Let k be a local field with finite residue field $\bar{k}$ of odd order. If $\mathbb{Q}_3 \subset k$ we suppose that $[k : \mathbb{Q}_3]$ is odd. Then there is a natural (split) short exact sequence of the form*

$$0 \rightarrow \mathcal{P}(\bar{k})[\frac{1}{2}]^{\oplus 2} \rightarrow H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \rightarrow K_3^{\mathrm{ind}}(k)[\frac{1}{2}] \oplus \mathcal{P}(k)[\frac{1}{2}] \rightarrow 0.$$

Example 8.4. In particular, if $p \geq 3$ is prime there is a natural decomposition

$$H_3(\mathrm{SL}_2(\mathbb{Q}_p[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \cong K_3^{\mathrm{ind}}(\mathbb{Q}_p)[\frac{1}{2}] \oplus \mathcal{P}(\mathbb{Q}_p)[\frac{1}{2}] \oplus \mathcal{P}(\mathbb{F}_p)[\frac{1}{2}]^{\oplus 2}.$$

We consider the case $k = \mathbb{Q}$. By Theorem 6.5 (6) there is a surjective map

$$\mathcal{RB}_0(\mathbb{Q})[\frac{1}{2}] \twoheadrightarrow \bigoplus_{p \text{ prime}} \mathcal{P}(\mathbb{F}_p)[\frac{1}{2}] \cong \bigoplus_p \mathbb{Z}/(p+1)'.$$

It is an open question whether this map is an isomorphism. We note, however, that Theorem 6.1 of [7] implies that $\mathcal{RB}_0(\mathbb{Q})$ is a torsion group.

Furthermore, we have $K_3^{\mathrm{ind}}(\mathbb{Q}) \cong \mathbb{Z}/24$ and $\mathcal{B}(\mathbb{Q}) \cong \mathbb{Z}/6$ so that

$$K_3^{\mathrm{ind}}(\mathbb{Q})[\frac{1}{2}] \cong \mathcal{B}(\mathbb{Q})[\frac{1}{2}] \cong \mathbb{Z}/3.$$

From the exact sequence

$$0 \rightarrow \mathcal{B}(\mathbb{Q}) \rightarrow \mathcal{P}(\mathbb{Q}) \rightarrow \mathcal{S}_2^2(\mathbb{Q}^\times) \rightarrow K_2(\mathbb{Q}) \rightarrow 0$$

and the fact that $K_2(\mathbb{Q})$ is a torsion group, we deduce that

$$\mathcal{P}(\mathbb{Q}) \left[\frac{1}{2} \right] \cong \mathbb{Z}/3 \oplus V$$

where V is a free $\mathbb{Z} \left[\frac{1}{2} \right]$ -module of countably infinite rank. Thus by Proposition 8.1 there is a short exact sequence

$$0 \rightarrow \mathcal{RB}_0(\mathbb{Q}) \left[\frac{1}{2} \right]^{\oplus 2} \rightarrow H_3(\mathrm{SL}_2(\mathbb{Q}[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow \mathbb{Z}/3 \oplus \mathbb{Z}/3 \oplus V \rightarrow 0.$$

Finally, we use the results of section 7 to calculate the kernel and cokernel of the stabilization map from $H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right])$ to $H_3(\mathrm{SL}(k[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right])$:

Theorem 8.5. *Let k be an infinite field. The cokernel of the map*

$$H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow H_3(\mathrm{SL}(k[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right])$$

is naturally isomorphic to $K_3^M(k) \left[\frac{1}{2} \right] \oplus K_2^M(k) \left[\frac{1}{2} \right]$.

The kernel of this map is naturally isomorphic to $\mathcal{RB}_0(k) \left[\frac{1}{2} \right] \oplus \mathcal{RP}_1(k) \left[\frac{1}{2} \right]$.

Proof. For any ring A , Suslin has shown ([19]) that the Hurewicz homomorphism $K_3(A) \rightarrow H_3(\mathrm{GL}(A), \mathbb{Z})$ induces an isomorphism

$$\frac{K_3(A)}{\{-1\} \cdot K_2(A)} \cong H_3(\mathrm{SL}(A), \mathbb{Z}).$$

Since the element $\{-1\} \in K_1(A)$ has order 2, it follows that there is an induced isomorphism $K_3(A) \left[\frac{1}{2} \right] \cong H_3(\mathrm{SL}(A), \mathbb{Z} \left[\frac{1}{2} \right])$.

In particular, for an infinite field k

$$\begin{aligned} H_3(\mathrm{SL}k[t, t^{-1}], \mathbb{Z} \left[\frac{1}{2} \right]) &\cong K_3(k[t, t^{-1}]) \left[\frac{1}{2} \right] \\ &\cong K_3(k) \left[\frac{1}{2} \right] \oplus K_2(k) \left[\frac{1}{2} \right] \cong H_3(\mathrm{SL}(k), \mathbb{Z} \left[\frac{1}{2} \right]) \oplus H_2(\mathrm{SL}(k), \mathbb{Z} \left[\frac{1}{2} \right]). \end{aligned}$$

Thus the stabilization map induces an map of short exact sequences with compatible splittings

$$\begin{array}{ccccccc} 0 & \longrightarrow & H_3(\mathrm{SL}_2(k), \mathbb{Z} \left[\frac{1}{2} \right]) & \longrightarrow & H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right]) & \longrightarrow & \mathcal{RP}_1(k) \left[\frac{1}{2} \right] \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H_3(\mathrm{SL}(k), \mathbb{Z} \left[\frac{1}{2} \right]) & \longrightarrow & H_3(\mathrm{SL}(k[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right]) & \longrightarrow & K_2(k) \left[\frac{1}{2} \right] \longrightarrow 0 \end{array}$$

The kernel of the stabilization map $H_3(\mathrm{SL}_2(k), \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow H_3(\mathrm{SL}(k), \mathbb{Z} \left[\frac{1}{2} \right])$ is isomorphic to $\mathcal{RB}_0(k) \left[\frac{1}{2} \right]$ and the cokernel is isomorphic to $K_3^M(k) \left[\frac{1}{2} \right]$.

We conclude by showing that the map $\mathcal{RP}_1(k) \left[\frac{1}{2} \right] \rightarrow K_2(k) \left[\frac{1}{2} \right]$ in the diagram above is the zero map:

Recall that the map $\Delta : H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow \mathcal{RP}_1(k) \left[\frac{1}{2} \right]$ factors as

$$H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow H_3(\mathrm{SL}_2(k(t)), \mathbb{Z} \left[\frac{1}{2} \right]) \rightarrow \mathcal{RB}(k(t)) \left[\frac{1}{2} \right] \xrightarrow{\delta_t} \mathcal{RP}_1(k) \left[\frac{1}{2} \right]$$

and there is a compatible factorization of the map $H_3(\mathrm{SL}(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \rightarrow K_2(k)$:

$$H_3(\mathrm{SL}(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \rightarrow H_3(\mathrm{SL}(k(t)), \mathbb{Z}[\frac{1}{2}]) \cong K_3(k(t))[\frac{1}{2}] \xrightarrow{\delta_t} K_2(k)[\frac{1}{2}].$$

Let $\alpha \in \mathcal{RP}_1(k)[\frac{1}{2}]$. Consider the element $\langle\langle t \rangle\rangle \cdot \alpha \in \mathcal{RP}_1((k(t))[\frac{1}{2}])$. Note that $\delta_t(\langle\langle t \rangle\rangle \cdot \alpha) = \alpha$.

However

$$\langle\langle t \rangle\rangle \cdot \alpha \in \mathcal{RB}(k(t))[\frac{1}{2}] = \mathrm{Ker}(\mathcal{RP}_1(k(t))[\frac{1}{2}] \rightarrow S_{\mathbb{Z}}^2(k(t)^\times))$$

since $\langle t \rangle$ acts trivially on $S_{\mathbb{Z}}^2(k(t)^\times)$.

Likewise

$$\langle\langle t \rangle\rangle \cdot \alpha \in \mathcal{RB}_0(k(t))[\frac{1}{2}] = \mathrm{Ker}(\mathcal{RB}(k(t))[\frac{1}{2}] \rightarrow \mathcal{P}(k(t))[\frac{1}{2}])$$

since $\langle t \rangle$ acts trivially on $\mathcal{P}(k(t))$.

But

$$\mathcal{RB}_0(k(t))[\frac{1}{2}] \cong \mathrm{Ker}(H_3(\mathrm{SL}_2(k(t)), \mathbb{Z}[\frac{1}{2}]) \rightarrow H_3(\mathrm{SL}(k(t)), \mathbb{Z}[\frac{1}{2}]))$$

It follows that if $\beta \in H_3(\mathrm{SL}_2(k(t)), \mathbb{Z}[\frac{1}{2}])$ is any lifting of $\langle\langle t \rangle\rangle \cdot \alpha$, then β maps to 0 in $H_3(\mathrm{SL}(k(t)), \mathbb{Z}[\frac{1}{2}])$.

Therefore, α maps to 0 in $K_2(k)[\frac{1}{2}]$ as claimed. $\square$

Corollary 8.6. *Let k be a quadratically closed or a real-closed field. Then the kernel of the stabilization map*

$$H_3(\mathrm{SL}_2(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}]) \rightarrow H_3(\mathrm{SL}(k[t, t^{-1}]), \mathbb{Z}[\frac{1}{2}])$$

is naturally isomorphic to $\mathcal{P}(k)[\frac{1}{2}]$.

REFERENCES

- [1] P. M. Cohn. On the structure of the GL_2 of a ring. *Inst. Hautes Études Sci. Publ. Math.*, (30):5–53, 1966. 4
- [2] Johan L. Dupont and Chih Han Sah. Scissors congruences. II. *J. Pure Appl. Algebra*, 25(2):159–195, 1982. 13
- [3] Kevin Hutchinson. Scissors congruence groups and the third homology of SL_2 of local rings and fields. *arXiv:1309.5010*. 2, 16
- [4] Kevin Hutchinson. A refined Bloch group and the third homology of SL_2 of a field. *J. Pure Appl. Algebra*, 217:2003–2035, 2013. 2, 14, 15
- [5] Kevin Hutchinson. A Bloch-Wigner complex for SL_2 . *J. K-Theory*, 12(1):15–68, 2013. 13, 14, 17
- [6] Kevin Hutchinson and Liqun Tao. The third homology of the special linear group of a field. *J. Pure Appl. Algebra*, 213:1665–1680, 2009. 2, 14
- [7] Kevin P. Knudson. The homology of $\mathrm{SL}_2(F[t, t^{-1}])$. *J. Algebra*, 180(1):87–101, 1996. 1, 3, 21
- [8] Kevin P. Knudson. The homology of special linear groups over polynomial rings. *Ann. Sci. École Norm. Sup. (4)*, 30(3):385–415, 1997. 1, 2
- [9] Kevin P. Knudson. *Homology of linear groups*, volume 193 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2001. 3, 4
- [10] Sava Krstić and James McCool. Free quotients of $\mathrm{SL}_2(R[x])$. *Proc. Amer. Math. Soc.*, 125(6):1585–1588, 1997. 2
- [11] A. Mazzoleni. A new proof of a theorem of Suslin. *K-Theory*, 35(3-4):199–211 (2006), 2005. 6
- [12] Fabien Morel. An introduction to $\mathbb{A}^1$ -homotopy theory. In *Contemporary developments in algebraic K-theory*, ICTP Lect. Notes, XV, pages 357–441 (electronic). Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004. 4
- [13] Fabien Morel. Sur les puissances de l’idéal fondamental de l’anneau de Witt. *Comment. Math. Helv.*, 79(4):689–703, 2004. 5
- [14] Fabien Morel. $\mathbb{A}^1$ -algebraic topology over a field, volume 2052 of *Lecture Notes in Mathematics*. Springer, Heidelberg, 2012. 6
- [15] Walter Parry and Chih-Han Sah. Third homology of $\mathrm{SL}(2, \mathbf{R})$ made discrete. *J. Pure Appl. Algebra*, 30(2):181–209, 1983. 14

- [16] Daniel Quillen. Higher algebraic K -theory. I. In *Algebraic K-theory, I: Higher K-theories (Proc. Conf., Battelle Memorial Inst., Seattle, Wash., 1972)*, pages 85–147. Lecture Notes in Math., Vol. 341. Springer, Berlin, 1973. 1
- [17] Jean-Pierre Serre. *Trees*. Springer-Verlag, Berlin, 1980. Translated from the French by John Stillwell. 3
- [18] A. A. Suslin. Torsion in K_2 of fields. *K-Theory*, 1(1):5–29, 1987. 4
- [19] A. A. Suslin. K_3 of a field, and the Bloch group. *Trudy Mat. Inst. Steklov.*, 183:180–199, 229, 1990. Translated in Proc. Steklov Inst. Math. **1991**, no. 4, 217–239, Galois theory, rings, algebraic groups and their applications (Russian). 2, 13, 22
- [20] Charles A. Weibel. *The K-book*, volume 145 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2013. An introduction to algebraic K -theory. 12

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY COLLEGE DUBLIN

E-mail address: kevin.hutchinson@ucd.ie